arXiv:1211.7314v1 [math.GT] 30 Nov 2012

VOLUMES OF THE SETS OF TRANSLATION SURFACES WITH SMALL SADDLE CONNECTIONS

DUC-MANH NGUYEN

ABSTRACT. In this paper, we prove some estimates of the volumes of the sets of translation surfaces of area one which have several small, disjoint saddle connections. These estimates can be viewed as generalizations of a result of Masur-Smillie. The main idea of the proof is based on an upper bound of some integrals on the spaces of translation surfaces with marked saddle connections which already appeared in [N12].

1. INTRODUCTION

1.1. **Abelian differentials.** Fix $\underline{k} = (k_1, \dots, k_n)$, $k_i \in \mathbb{N}$, and let $\mathcal{H}(\underline{k})$ denote the moduli space of pairs (M, ω), where M is a compact Riemann surface, and ω is a holomorphic 1-form which has exactly n zeros with orders $k_1, \dots, k_n$. The one-form ω defines a flat metric structure on M whose transition maps are translations $z \mapsto z + c$ of $\mathbb{R}^2 \simeq \mathbb{C}$. The singularities of this metric are the zeros of ω, the cone angle at a zero of order k is $2\pi(k+1)$. The space $\mathcal{H}(\underline{k})$ is called a *stratum* of the moduli space of translation surfaces of genus g, where g and $\underline{k}$ are related by the equation

$$k_1 + \dots + k_n = 2g - 2.$$

We denote by $\mathcal{H}_1(\underline{k})$ be the subset of $\mathcal{H}(\underline{k})$ consisting of surfaces of area one (with respect to the flat metric). It is sometimes useful to consider translation surfaces with marked regular points, in this situation, we consider the marked points as zeros of order 0 of ω. This convention does not affect the arguments in what follows.

It is well known that the space $\mathcal{H}(\underline{k})$ is a complex algebraic orbifold, equipped with a natural volume form μ coming from the period mapping (see Section 2.2). We have $\dim_{\mathbb{C}} \mathcal{H}(\underline{k}) = 2g + n - 1$. Let μ_1 be the volume form on $\mathcal{H}_1(\underline{k})$ which is defined by the formula

$$d\mu = d\mu_1 d\mathrm{A},$$

where A is the area function.

Given $(M, \omega) \in \mathcal{H}(\underline{k})$, a *saddle connection* on M is a geodesic segment for the flat metric whose endpoints are singularities (the endpoints are not necessarily distinct). We denote the length of a saddle connection γ by $|\gamma|$.

Definition 1.1. *Let $\{\gamma_1, \dots, \gamma_m\}$ be a family of saddle connections on M. This family will be called an* independent family *if the following conditions are fulfilled*

(i) $\mathrm{int}(\gamma_i) \cap \mathrm{int}(\gamma_j) = \varnothing$ *if* $i \neq j$,

(ii) *the family* $\{[\gamma_1], \dots, [\gamma_m]\}$ *is linearly independent in* $H_1(M, \{p_1, \dots, p_n\}; \mathbb{R})$, *where* $p_1, \dots, p_n$ *are zeros of* ω.

Given $\underline{\epsilon} = (\epsilon_1, \dots, \epsilon_m) \in (\mathbb{R}_{>0})^m$, let us de note by $\mathcal{H}^{(m)}_{1,\underline{\epsilon}}(\underline{k})$ the subset of $\mathcal{H}_1(\underline{k})$ consisting of surfaces $(M, \omega) \in \mathcal{H}_1(\underline{k})$ on which there exists an ordered independent family of saddle connections $\{\gamma_1, \dots, \gamma_m\}$ such that

$$|\gamma_j| < \epsilon_j, \ \ j = 1, \dots, m.$$

When ϵ_j are small enough, such a family only exists if $m < \dim_{\mathbb{C}} \mathcal{H}(\underline{k}) = 2g + n - 1$. In this paper, we will prove the following theorem, which provides an estimate of the volume of the set $\mathcal{H}^{(m)}_{1,\underline{\epsilon}}(\underline{k})$

Theorem 1.2. *For any* $m < \dim_{\mathbb{C}} \mathcal{H}(\underline{k})$, *there exists a constant* $C = C(\underline{k}, m)$ *such that*

$$\mu_1(\mathcal{H}^{(m)}_{1,\underline{\epsilon}}(\underline{k})) < C\epsilon_1^2 \dots \epsilon_m^2.$$

1.2. **Quadratic differentials.** Let M be a compact Riemann surface, and ϕ be a meromorphic quadratic differential on M. The quadratic deferential ϕ defines on M a flat metric structure whose transition maps are of the form $z \mapsto \pm z + c, c$ is constant. If $\int_M |\phi|^2$ is finite, which implies that all the poles of ϕ are simple, the area of this metric structure is finite, and its singularities are isolated cone points corresponding to the zeros and poles of ϕ. The cone angle at a zero of order k is $(k + 2)\pi$, where a zero of order -1 is a simple pole.

In the case $\phi = \omega^2$, where ω is a holomorphic one-form, the flat metrics defined by ϕ and ω are the same, thus we will be only interested in quadratic differentials which are not square of any holomorphic one-form. In this case, the corresponding flat surface is sometimes called a half-translation surface.

There is a natural stratification on the set of quadratic differentials by the orders of their zeros, we denote by $\mathcal{Q}(\underline{d})$, $\underline{d} = (d_1, \dots, d_n), d_i \in \{-1, 0, 1, 2, \dots\}$, the set of pairs (M, ϕ) where ϕ has exactly n zeros, with orders given by $\underline{d}$, and ϕ is not the square of a holomorphic one-form. By convention, a zero of order 0 is a marked regular point of the flat metric. Remark that the genus of M is determined by the formula

$$d_1 + \dots + d_n = 4g - 4.$$

It is well known that $\mathcal{Q}(\underline{d})$ is also a complex algebraic orbifold of dimension $2g + n - 2$, which is equipped with a natural Lebesgue volume form μ. Let $\mathcal{Q}_1(\underline{d})$ denote the subset of $\mathcal{Q}(\underline{d})$ consisting of surface of area one, then we have a volume form μ_1 on $\mathcal{Q}_1(\underline{d})$ which is defined by

$$d\mu = d\mathrm{A} d\mu_1,$$

where A is the area function on $\mathcal{Q}(\underline{d})$.

Given a meromorphic quadratic differential (M, ϕ), which is not a square of a holomorphic one-form, there exists a canonical (ramified) double covering $\pi : \hat{M} \to M$ such that the quadratic differential $\hat{\phi} = \pi^*(\phi)$ is the square of a holomorphic one-form $\hat{\omega}$ on $\hat{M}$. We have an involution $\tau : \hat{M} \to \hat{M}$ satisfying $q = \tau(p)$ if and only if $\pi(p) = \pi(q)$, and $\tau^*\hat{\omega} = -\hat{\omega}$. Let S denote the set of singularities of the flat metric structure defined by ϕ on M. We will call a geodesic segment γ with endpoints in S a *saddle connection* of M. The pre-image of γ in $\hat{M}$ consists of two geodesic segments $\hat{\gamma}^1, \hat{\gamma}^2$ with endpoints in $\hat{S} = \pi^{-1}(S)$. We choose the orientation of $\hat{\gamma}^1$ and $\hat{\gamma}^2$ so that $\tau(\hat{\gamma}^1) = -\hat{\gamma}^2$ and $\tau(\hat{\gamma}^2) = -\hat{\gamma}^1$. We will denote by $[\hat{\gamma}]$ the homology class in $H_1(\hat{M}, \hat{S}; \mathbb{Z})$ represented by $[\hat{\gamma}^1] + [\hat{\gamma}^2]$. We have a natural decomposition

$$H_1(\hat{M}, \hat{S}; \mathbb{R}) = H_1(\hat{M}, \hat{S}; \mathbb{R})^- \oplus H_1(\hat{M}, \hat{S}; \mathbb{R})^+,$$

where $H_1(\hat{M}, \hat{S}; \mathbb{R})^\pm$ are the eigenspaces of τ corresponding to the eigenvalues ± 1 respectively. By definition, $[\hat{\gamma}]$ belongs to $H_1(\hat{M}, \hat{S}; \mathbb{R})^-$, and $\hat{\omega}$ belongs to $H^1(\hat{M}, \hat{S}; \mathbb{C})^- = \mathrm{Hom}_{\mathbb{R}}(H_1(\hat{M}, \hat{S}; \mathbb{R})^-, \mathbb{C})$.

Definition 1.3. *Let* $\{\gamma_1, \dots, \gamma_m\}$ *be a family of saddle connections on* M. *We will say that* $\{\gamma_1, \dots, \gamma_m\}$ *is an* independent family *if the following conditions are fulfilled*

(i) $\mathrm{int}(\gamma_i) \cap \mathrm{int}(\gamma_j) =$ *if* $i \neq j$,

(ii) *the family* $\{[\hat{\gamma}_1], \dots, [\hat{\gamma}_m]\}$ *is independent in* $H_1(\hat{M}, \hat{S}; \mathbb{R})^-$.

For all $\underline{\epsilon} = (\epsilon_1, \dots, \epsilon_m) \in (\mathbb{R}_{>0})^m$, where $m < \dim_{\mathbb{C}} \mathcal{Q}(\underline{d}) = 2g + n - 2$, let us denote by $\mathcal{Q}^{(m)}_{1,\underline{\epsilon}}(\underline{d})$ the set of $(M, \phi) \in \mathcal{Q}_1(\underline{d})$ such that there exists an ordered independent family of saddle connections $\{\gamma_1, \dots, \gamma_m\}$ on M satisfying

$$|\gamma_j| < \epsilon_j, \ j = 1, \dots, m.$$

Similar to the case of Abelian differentials, we will prove

Theorem 1.4. *For any* $m < \dim_{\mathbb{C}} \mathcal{Q}(\underline{d})$, *there exists a constant* $C = C(\underline{d}, m)$ *such that for every* $\underline{\epsilon} \in (\mathbb{R}_{>0})^m$, *we have*

$$\mu_1(\mathcal{Q}^{(m)}_{1,\underline{\epsilon}}(\underline{d})) < C\epsilon_1^2 \dots \epsilon_m^2.$$

1.3. **Outline.** The main results of this paper are proven as consequences of some estimates of integrals over the spaces of (half-)translation surfaces with marked saddle connections. Roughly speaking, for each $\underline{\epsilon}$ we will define a function $\mathfrak{F}_{\underline{\epsilon}}$ on the spaces of (half-)translation surfaces with m-marked saddle connections which involves the area of the surface and the lengths of the marked saddle connections. Theorems 1.2 and 1.4 then follow from the fact that the integral of $\mathfrak{F}_{\underline{\epsilon}}$ is bounded by $K\epsilon_1^2 \dots \epsilon_m^2$ (see Theorems 2.3 and 6.1), where K is a constant. The proof of these estimates uses construction of translation surfaces from triangles in $\mathbb{R}^2$, and similar ideas as in [N12].

This paper is inspired from the work [EMZ03, MZ08] of Eskin, Masur, and Zorich in which they carefully study the neighborhood of the boundary of moduli spaces of Abelian differentials and quadratic differentials to obtain formulae relating the Siegel-Veech constants and the volumes of the strata. One important ingredient of the proof of their results is a result of Masur-Smillie, which gives an estimate on the volume of the set of quadratic differentials which have two small non-parallel saddle connections (see [EMZ03], Lemma 7.1 and [MS91] Theorem 10.3). In this perspective, Theorem 1.2 and Theorem 1.4 can be considered as a generalization of Masur-Smillie's estimate.

2. TRANSLATION SURFACE WITH MARKED SADDLE CONNECTIONS

2.1. Translation surfaces with marked saddle connections.

Definition 2.1. *Let us denote by* $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ *the moduli space of triples* $(M,\omega,\{\gamma_1,\dots,\gamma_m\})$, *where*

- (M,ω) *belongs to* $\mathcal{H}(\underline{k})$,
- $\{\gamma_1,\dots,\gamma_m\}$ *is an independent ordered family of saddle connections.*

Let $X=(M,\omega,\{\gamma_1,\dots,\gamma_m\})$ and $X'=(M',\omega',\{\gamma'_1,\dots,\gamma'_m\})$ be two elements in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$. We say that X and X' have the same *topological type* if there exists a homeomorphism $\varphi: M\to M'$ which realizes a bijection between the sets of zeros of ω and ω' preserving the orders, such that $\varphi(\gamma_i)=\gamma'_i$. For a fixed vector $\underline{k}=(k_1,\dots,k_n)$, there are only finitely many topological types for surfaces in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$.

2.2. Local chart and volume form. Let us first recall the definitions of local charts and volume form on $\mathcal{H}(\underline{k})$ using the period mapping. Let (M,ω) be a pair in $\mathcal{H}(\underline{k})$. The zeros of ω are denoted by $p_1,\dots,p_n$, and their orders by is k_i respectively. Let $\{c_1,\dots,c_{2g+n-1}\}$ be a set of curves on M which is a basis of $H_1(M,\{p_1,\dots,p_n\};\mathbb{Z})$. For any element (M',ω') close enough to (M,ω) in $\mathcal{H}(\underline{k})$, we denote by $\{c'_1,\dots,c'_{2g+n-1}\}$ the corresponding curves on M'. We can then define a map Φ from a neighborhood of (M,ω) into $\mathbb{C}^{2g+n-1}$, which sends a pair (M',ω') to the vector $(\int_{c'_1}\omega',\dots,\int_{c'_{2g+n-1}}\omega')$. The map Φ is called the *period mapping*. It is well known that Φ is a local chart for $\mathcal{H}(\underline{k})$, hence $\mathcal{H}(\underline{k})$ has a complex orbifold structure of dimension $2g+n-1$.

Let $\lambda_{2(2g+n-1)}$ denote the Lebesgue measure of $\mathbb{C}^{2g+n-1}\simeq\mathbb{R}^{2(2g+n-1)}$. Since the bases of $H_1(M,\{p_1,\dots,p_n\};\mathbb{Z})$ are related by elements of the group $\mathrm{GL}(2g+n-1,\mathbb{Z})$, the volume form $\Phi^*\lambda_{2(2g+n-1)}$ is well defined on $\mathcal{H}(\underline{k})$. We denote this volume form by μ.

Let $\mathcal{H}_1(\underline{k})$ denote the subspace of $\mathcal{H}(\underline{k})$ consisting of pairs (M,ω) such that $\int_M|\omega|^2=1$. An element of $\mathcal{H}_1(\underline{k})$ corresponds to a translation surface of area 1. The volume form μ induces a volume form μ_1 on $\mathcal{H}_1(\underline{k})$ by the following relation

$$d\mu = d\mathrm{A}d\mu_1.$$

where A is the area function. It is proved by Masur [M82] and Veech [V90] that the total volume of $\mathcal{H}_1(\underline{k})$ with respect to μ_1 is finite.

There is an obvious projection $\varrho:\widetilde{\mathcal{H}}^{(m)}(\underline{k})\to\mathcal{H}(\underline{k})$, which maps the triple $(M,\omega,\{\gamma_1,\dots,\gamma_m\})$ to the pair (M,ω). If $(M',\omega')\in\mathcal{H}(\underline{k})$ is close enough to (M,ω), then there also exists m saddle connections $\gamma'_1,\dots,\gamma'_m$ on M' such that $X:=(M,\omega,\{\gamma_1,\dots,\gamma_m\})$ and $X':=(M',\omega',\{\gamma'_1,\dots,\gamma'_m\})$

have the same topological type. Therefore, we can identify a neighborhood of X in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ with a neighborhood of (M,ω) in $\mathcal{H}(\underline{k})$, and use Φ to define local charts for $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$. Since $\{[\gamma_1],\dots,[\gamma_m]\}$ is an independent family in $H_1(M,\{p_1,\dots,p_n\};\mathbb{Z})$, we can choose a basis $\{c_1,\dots,c_{2g+n-1}\}$ of $H_1(M,\{p_1,\dots,p_n\};\mathbb{Z})$ such that $c_1=\gamma_1,\dots,c_m=\gamma_m$.

The projection ϱ is clearly locally homeomorphic. Thus, we can pullback the volume form μ of $\mathcal{H}(\underline{k})$ to $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$. By a slight abuse of notation, we will also denote by μ this volume form on $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$. Let us now resume what we have seen in the following

Theorem 2.2. $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ *admits a flat complex affine orbifold structure (that is, the transition maps are complex linear transformations) of dimension* $2g+n-1$, *for which* μ *is a parallel volume form (*i.e. *invariant by transition maps of the flat complex affine structure).*

2.3. **Energy functions.** For every $\underline{\epsilon}=(\epsilon_1,\dots,\epsilon_m)$, with $\epsilon_j>0$, we define a function $\mathfrak{F}_{\underline{\epsilon}}:\widetilde{\mathcal{H}}^{(m)}(\underline{k})\to\mathbb{R}$ by the following formula

$$\mathfrak{F}_{\underline{\epsilon}}:(M,\omega,\{\gamma_1,\dots,\gamma_m\})\mapsto \exp\Big(-\sum_{j=1}^{m}\frac{|\gamma_j|^2}{\epsilon_j^2}-\mathrm{A}(M)\Big).$$

Theorem 1.2 is a consequence of the following

Theorem 2.3. *There exists a constant* $K=K(\underline{k},m)$ *such that*

$$\int_{\widetilde{\mathcal{H}}^{(m)}(\underline{k})}\mathfrak{F}_{\underline{\epsilon}}d\mu<K\epsilon_1^2\dots\epsilon_m^2.$$

The proof of this theorem will be given in Section 5

2.4. **Proof of Theorem 1.2.**

Proof. Let us now give the proof of 1.2 using Theorem 2.3. Let $\mathcal{H}^{(m)}_{\underline{\epsilon}}(\underline{k})$ be the subset of $\mathcal{H}(\underline{k})$ consisting of surfaces (M,ω) on which there exists an independent family of saddle connections $\{\gamma_1,\dots,\gamma_m\}$ such that

$$|\gamma_j|^2<\epsilon_j^2\mathrm{A}(M).$$

Note that $\mathcal{H}^{(m)}_{\underline{\epsilon}}(\underline{k})=\mathbb{R}_+^*\cdot\mathcal{H}^{(m)}_{1,\underline{\epsilon}}(\underline{k})$, where the action of $\mathbb{R}_+^*$ is given by $t\cdot(M,\omega)=(M,t\omega),\ t\in\mathbb{R}_+^*$. Let $\widetilde{\mathcal{H}}^{(m)}_{\underline{\epsilon}}(\underline{k})\subset\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ be the pre-image of $\mathcal{H}^{(m)}_{\underline{\epsilon}}(\underline{k})$ by the natural projection $\varrho:\widetilde{\mathcal{H}}^{(m)}(\underline{k})\to\mathcal{H}(\underline{k})$. Theorem 2.3 clearly implies

$$\int_{\widetilde{\mathcal{H}}^{(m)}_{\underline{\epsilon}}(\underline{k})}\mathfrak{F}_{\underline{\epsilon}}d\mu<K\epsilon_1^2\dots\epsilon_m^2. \tag{1}$$

Let us define the function $f_{\underline{\epsilon}}:\mathcal{H}(\underline{k})\to\mathbb{R}$ as follows

$$f_{\underline{\epsilon}} : (M,\omega) \mapsto e^{-\mathrm{A}(M)} \left(\sum_{\substack{\{\gamma_1,\dots,\gamma_m\} \\ \text{independent family in } M}} e^{-(\frac{|\gamma_1|^2}{\epsilon_1^2}+\dots+\frac{|\gamma_m|^2}{\epsilon_m^2})} \right).$$

By definition, we have

$$\int_{\widetilde{\mathcal{H}}_{\underline{\epsilon}}^{(m)}(\underline{k})} \mathfrak{F}_{\underline{\epsilon}} d\mu = \int_{\mathcal{H}_{\underline{\epsilon}}^{(m)}(\underline{k})} f_{\underline{\epsilon}} d\mu.$$

But if $(M,\omega) \in \mathcal{H}_{\underline{\epsilon}}^{(m)}(\underline{k})$, then by definition there exists at least an independent family $\{\gamma_1,\dots,\gamma_m\}$ in M such that

$$e^{-(\frac{|\gamma_1|^2}{\epsilon_1^2}+\dots+\frac{|\gamma_m|^2}{\epsilon_m^2})} > e^{-m\mathrm{A}(M)}.$$

Thus on $\mathcal{H}_{\underline{\epsilon}}^{(m)}(\underline{k})$, we have

$$f_{\underline{\epsilon}}(.) > e^{-(m+1)\mathrm{A}(.)},$$

and (1) implies

$$\int_{\mathcal{H}_{\underline{\epsilon}}^{(m)}(\underline{k})} e^{-(m+1)\mathrm{A}} d\mu < K\epsilon_1^2\dots\epsilon_m^2. \tag{2}$$

Recall that the space $\mathcal{H}(\underline{k})$ can be locally identify with an open of $\mathbb{R}^{2N}, N = 2g+n-1$, so that μ is the Lebesgue measure. By this identification, $\mathcal{H}_1(\underline{k})$ corresponds to the hypersurface $\mathbf{Q}_1$ defined by the equation $\mathrm{A} = 1$. Let us make the following change of coordinates

$$\begin{array}{rccc} \Phi : & \mathbf{Q}_1\times]0,+\infty[& \rightarrow & \mathbb{R}^{2N} \\ & (v,t) & \mapsto & t\cdot v. \end{array}$$

Note that we have $\Phi^*\mathrm{A}(v,t) = t^2$. Using the relation $d\mu = d\mathrm{A}d\mu_1$ on $\mathbf{Q}_1$, one can easily check that

$$\Phi^* d\mu = 2t^{2N-1}dt d\mu_1.$$

Therefore,

$$\int_{\mathcal{H}_{\underline{\epsilon}}^{(m)}(\underline{k})} e^{-(m+1)\mathrm{A}} d\mu = \int_{\mathcal{H}_{1,\underline{\epsilon}}^{(m)}(\underline{k})} \left(\int_0^{+\infty} 2t^{2N-1} e^{-(m+1)t^2} dt \right) d\mu_1.$$

It follows immediately from (2) that

$$\mu_1(\mathcal{H}^{(m)}_{1,\underline{\epsilon}}(\underline{k})) < C\epsilon_1^2\dots\epsilon_m^2.$$

Theorem 1.2 is then proven.

□

3. TRIANGULATIONS AND LOCAL CHARTS

Let $X := (M, \omega, \{\gamma_1, \dots, \gamma_m\})$ be a point in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$. It is a well known fact that there always exists a geodesic triangulation T of M whose vertices are the singular points such that the family $\{\gamma_1, \dots, \gamma_m\}$ is included in the 1-skeleton $\mathrm{T}^{(1)}$ of T. We will call such a triangulation an *admissible triangulation*. Let N_1, N_2 denote the number of edges and the number of triangles of T respectively. Note that N_1 and N_2 are completely determined by the genus g of M and the number n of zeros of ω, namely, we have

$$N_1 = 3(2g+n-2) \text{ and } N_2 = 2(2g+n-2).$$

Choosing for each edge of T an orientation, we can then associate to each oriented edge e a complex number $Z(e) = \int_e \omega$. Thus, the triangulation T provides us with a vector $Z \in \mathbb{C}^{N_1}$. The coordinates of Z must satisfy the following condition: if e_i, e_j, e_k are three edges of a triangle of T, then we have

$$\pm Z(e_i) \pm Z(e_j) \pm Z(e_k) = 0. \tag{3}$$

The signs of $Z(e_i), Z(e_j), Z(e_k)$ depend on their orientation. We have N_2 equations of type (3), each of which corresponds to a triangle in $\mathrm{T}^{(2)}$. Let $\mathbf{S}_{\mathrm{T}}$ denote the system consisting of those linear equations, and V_{T} be the subspace of solutions of $\mathbf{S}_{\mathrm{T}}$ in $\mathbb{C}^{N_1}$.

Lemma 3.1. *Let Z' be a vector in V_{T}. If Z' is close enough to Z then there exists*

- *an element $X' = (M', \omega', \{\gamma'_1, \dots, \gamma'_m\}) \in \widetilde{\mathcal{H}}^{(m)}(\underline{k})$ close to X,*
- *an admissible triangulation T' of M',*
- *a homeomorphism $\varphi : M \to M'$ which maps T to T',*

such that Z' is the vector associated to T'.

Proof. We consider Z as a function from $\mathrm{T}^{(1)}$ to $\mathbb{C}$ (where $\mathrm{T}^{(k)}$ is the k-skeleton of T, $k = 0, 1, 2$). We can construct a flat surface from $Z' \in \mathrm{V}_{\mathrm{T}}$ as follows

(a) construct a triangulation from $Z'(e_i), Z'(e_j), Z'(e_k)$, whenever there exists a triangle in $\mathrm{T}^{(2)}$ which contains e_i, e_j, e_k ((e_i, e_j, e_k) are distinct edges),
(b) glue two triangles together if the corresponding triangles in $\mathrm{T}^{(2)}$ have a common edge.

This construction provides us with a surface M' equipped with a flat metric structure with cone singularities. Since we only identify edges which have the same direction and the same length, the 1-form dz on $\mathbb{C}$ is preserved by the gluings. Therefore, we also get a holomorphic 1-form ω' on M', and hence a translation surface. As all the topological invariants (orientation, genus, number of singularities, cone angles) are discrete, it follows that if Z' is close enough to Z then (M',ω') has the same topological invariants as (M,ω), hence $(M',\omega')\in\mathcal{H}(\underline{k})$.

The surface (M',ω') comes equipped with a geodesic triangulation T', the triangles of T' are formed by triples of vectors $Z'(e_i), Z'(e_j), Z'(e_k)$ which satisfy an equation of type (3). From the construction of M', we have a homeomorphism $\varphi : M\rightarrow M'$ which sends a triangle of T to a triangle of T'. Set $\gamma'_j=\varphi(\gamma_j)$, then $(M',\omega',\{\gamma'_1,\dots,\gamma'_m\})$ is an element of $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ which has the same topological type as $(M,\omega,\{\gamma_1,\dots,\gamma_m\})$. ◻

From Lemma 3.1, we see that one can define a map

$$\Psi_{\mathrm{T}}:\mathcal{U}\rightarrow\widetilde{\mathcal{H}}^{(m)}(\underline{k}),$$

where $\mathcal{U}$ is an open subset of $\mathrm{V_T}$ that contains Z. Actually, we have

Proposition 3.2. *For $\mathcal{U}$ small enough, Ψ_{T} is continuous and injective, it realizes an homeomorphism between $\mathcal{U}$ and its image.*

Proof. Let us first show that $\dim_{\mathbb{C}}\mathrm{V_T}=2g+n-1$. Since $\dim H_1(M,\{p_1,\dots,p_n\};\mathbb{Z})=2g+n-1$, where $p_1,\dots,p_n$ are the zeros of ω, there always exists in $\mathrm{T}^{(1)}$ a family of $2g+n-1$ edges such that if we cut M along this family the resulting surface is a union of topological disks. Let $\mathcal{B}=\{\tilde{e}_1,\dots,\tilde{e}_{2g+n-1}\}$ be such a family.

Recall that every equation in $\mathbf{S}_{\mathrm{T}}$ corresponds to a triangle in $\mathrm{T}^{(2)}$, we write the equation associated to a triangle coherently with the orientation of its boundary. Since every edge e of T belongs to two triangles, $Z(e)$ appears in exactly two equations with opposite signs. Thus, the sum of all equations in $\mathbf{S}_{\mathrm{T}}$ is the trivial equation $0=0$, which implies that $\mathrm{rk}(\mathbf{S}_{\mathrm{T}})\leq N_2-1$, therefore we have

$$\dim\mathrm{V_T}\geq N_1-(N_2-1)=2g+n-1. \tag{4}$$

Assume that $e_1,\dots,e_k$ are the k edges of T which bound an open embedded disk D in M. The disk D is divided into triangles in $\mathrm{T}^{(2)}$, by taking the sum of the equations corresponding to the triangles contained in D, we get

$$\pm Z(e_1)\pm\cdots\pm Z(e_k)=0. \tag{5}$$

If $e\in\mathrm{T}^{(1)}$ is an edge which does not belong to the family $\mathcal{B}$, then e and some edges in $\mathcal{B}$ bound an embedded disk in M. It follows from (5) that we can write $Z(e)$ as a linear function of $(Z(\tilde{e}_1),\dots,Z(\tilde{e}_{2g+n-1}))$. Therefore, we have

$$\dim_{\mathbb{C}}\mathrm{V_T}\leq 2g+n-1. \tag{6}$$

From (4) and (6), we can conclude that $\dim_{\mathbb{C}}\mathrm{V_T}=2g+n-1$, and the map

$$\begin{array}{llll} \mathrm{F}_{\mathrm{T}}: & \mathrm{V}_{\mathrm{T}} & \rightarrow & \mathbb{C}^{2g+n-1} \\ & Z & \mapsto & (Z(\tilde{e}_1), \dots, Z(\tilde{e}_{2g+n-1})) \end{array}$$

is an isomorphism.

There exists a neighborhood $\mathcal{V}$ of $(M, \omega, \{\gamma_1, \dots, \gamma_m\})$ in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$, such that the period mapping $\Phi : (M, \omega) \mapsto (\int_{\tilde{e}_1} \omega, \dots, \int_{\tilde{e}_{2g+n-1}} \omega)$ is a local chart on $\mathcal{V}$. Consider the map

$$\Phi \circ \Psi_{\mathrm{T}} : \Psi_{\mathrm{T}}^{-1}(\mathcal{V}) \rightarrow \mathbb{C}^{2g+n-1}.$$

By definition, we see that $\Phi \circ \Psi_{\mathrm{T}}(Z) = \mathrm{F}_{\mathrm{T}}(Z) = (Z(\tilde{e}_1), \dots, Z(\tilde{e}_{2g+n-1}))$, therefore $\Phi \circ \Psi_{\mathrm{T}}$ is a restriction of the map F_{T}. Since F_{T} is an isomorphism, $\Phi \circ \Psi_{\mathrm{T}}$ is continuous and injective, and the proposition follows.

□

4. SPECIAL TRIANGULATION

4.1. **Construction.** From Proposition 3.2, we know that there is a locally homeomorphic map from some open subset of V_{T} into $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$. Our goal in this section is to specify an open subset of V_{T} on which this map is one-to-one. For this purpose, we first construct for each surface in an open dense set of $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ a particular triangulation which will be called *special triangulation*.

Let $X := (M, \omega, \{\gamma_1, \dots, \gamma_m\})$ be a point in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$, and $p_1, \dots, p_n$ be the zeros of ω. Let ξ be the vector field which is defined on $M \setminus \{p_1, \dots, p_n\}$ by the following condition, in any local chart of M such that $\omega = dz$, we have $\xi = \partial_y$. The orbits of the flow generated by ξ are called *vertical geodesics* in M.

Let $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$ denote the set of $(\Sigma, \xi, \{\gamma_1, \dots, \gamma_m\})$ verifying

(a) none of the connections $\{\gamma_1, \dots, \gamma_m\}$ is parallel to the field ξ,
(b) every geodesic ray emanating from the singular points of M, which are called *separatrices*, in directions $\pm\xi$ intersects the interior of one of the saddle connections $\{\gamma_1, \dots, \gamma_m\}$.

Lemma 4.1. *$\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$ is an open dense subset of $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$.*

Proof. It is not difficult to see that both conditions (a) and (b) are open conditions, thus $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$ is an open in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$.

To see that $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$ is dense, let us consider an element $X = (M, \omega, \{\gamma_1, \dots, \gamma_m\}) \in \widetilde{\mathcal{H}}^{(m)}(\underline{k})$. By a classical result (see for example [MT02]), we know that the set of $\theta \in \mathbb{S}^1$ such that the vertical flow on the surface $(M, e^{\imath\theta}\omega)$ is not minimal is countable, which implies that the set Θ of $\theta \in \mathbb{S}^1$ such that the flow in direction θ is minimal is dense in $\mathbb{S}^1$. For any $\theta \in \Theta$, if γ is a saddle connection, then the direction of γ is different from θ, and every separatrix in direction θ must intersect γ (since such a separatrix is dense in M). It follows in particular that we can find a sequence $\{\theta_k\} \subset \Theta$ converging to 0 such that $(M, e^{\imath\theta_k}\omega, \{\gamma_1, \dots, \gamma_m\})$ belongs to $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$. Therefore X is contained in the closure of $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$, and the lemma follows. □

Suppose from now on that X belongs to $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$. We truncate all the geodesic rays from the singularities of Σ in directions $\pm\xi$ at their first intersection with the set $\sqcup_{i=1}^m \mathrm{int}(\gamma_i)$. Note that, for any geodesic segment σ in Σ which is not parallel to ξ, by using the isometric embedding of a neighborhood of σ into $\mathbb{R}^2$ so that ξ becomes the vertical field $\partial_y = (0,1)$, we can specify the left and right endpoints of σ. In what follows, all the geodesic segments are oriented from the left to the right.

Pick a saddle connection γ in the family $\{\gamma_1,\dots,\gamma_m\}$. If η is a vertical geodesic ray from a singular point that intersects $\mathrm{int}(\gamma)$, we say that η is an upper ray (resp. a lower ray) at γ if η reaches γ from the upper side (resp. lower side). Equivalently, η is a upper side (resp. lower side) at γ if it starts from a singular point in direction $-\xi$ (resp. ξ).
Let A and B denote the left and right endpoints of γ respectively. We denote by $\eta_1^+,\dots,\eta_k^+$ the upper rays at γ from the left to the right. The endpoints of η_j^+ will be denoted by P_j and Q_j, where $Q_j \in \mathrm{int}(\gamma)$, and P_j is the singularity where the ray η_j^+ starts. Note that we may have $P_j = P_{j'}$ even if $j' \neq j$.

We will find a family of embedded triangles in M with disjoint interior whose union contains all the segments η_j^+ by the following procedure: let j_0 be the smallest index such that $|\eta_{j_0}^+| = \min\{|\eta_j^+|,\ j=1,\dots,k\}$. Using the developing map, we can realize γ and $\eta_{j_0}^+$ as two segments $\tilde{A}\tilde{B}$ and $\tilde{P}_{j_0}\tilde{Q}_{j_0}$ respectively in the plane $\mathbb{R}^2$, where $\tilde{Q}_{j_0} \in \tilde{A}\tilde{B}$, and $\tilde{P}_{j_0}\tilde{Q}_{j_0}$ is vertical. Let $\tilde{\Delta}$ denote the triangle in $\mathbb{R}^2$ with vertices $\tilde{A}, \tilde{B}, \tilde{P}_{j_0}$.

Claim 4.2. *There exists a locally isometric map* $\varphi : \tilde{\Delta} \to M$ *which satisfies:*

- $\varphi(\tilde{A}) = A, \varphi(\tilde{B}) = B, \varphi(\tilde{P}_{j_0}) = P_{j_0}$,
- $\varphi(\mathrm{int}(\tilde{\Delta}))$ *is disjoint from the family* $\{\gamma_1,\dots,\gamma_m\}$,
- *the restriction of* φ *to* $\mathrm{int}(\tilde{\Delta})$ *is an embedding.*

In particular, $\Delta = \varphi(\tilde{\Delta})$ *is an embedded geodesic triangle in M whose vertices are singularities of the flat metric.*

Proof. Let (ψ_t), $t \in \mathbb{R}$, denote the flow generated by ξ on M. Given a point p in $M \setminus \{p_1,\dots,p_n\}$, if there exists $s > 0$ such that $\psi_s(p) \in (\cup_{i=1}^m \gamma_i) \cup \{p_1,\dots,p_n\}$, then for every $t > s$, we consider by convention that $\psi_t(p) = \psi_s(p)$.

Set $t_0 = |\eta_{j_0}^+|$, and consider the set $L = \cup_{0\le t\le t_0}\psi_t(\gamma) \subset M$. Note that $P_{j_0} \in \psi_{t_0}(\gamma)$. If $\psi_t(\gamma)$ is disjoint from $\cup_{j=1}^m \gamma_j$ for all $t \in]0,t_0[$, then L is isometric to a parallelogram P in $\mathbb{R}^2$ with two vertical sides of length t_0, the lower side of this parallelogram is $\tilde{A}\tilde{B}$, and the upper side contains $\tilde{P}_{j_0}$. In general, L is isometric to a polygon $\tilde{L}$ included in the parallelogram P. The additional sides of $\tilde{L}$ correspond to subsegments of $\{\gamma_1,\dots,\gamma_m\}$. The hypothesis $|\eta_{j_0}^+| = \min\{\eta_j^+,\ j=1,\dots,k\}$ implies that $\tilde{L}$ is convex. Therefore, $\tilde{\Delta}$ is included in $\tilde{L}$.

We have naturally a map $\varphi : \tilde{\Delta} \to M$ which is locally isometric such that $\varphi(\tilde{A}) = A, \varphi(\tilde{B}) = B, \varphi(\tilde{P}_{j_0}) = P_{j_0}$. Since the polygon $\tilde{L}$ is convex, $\varphi(\mathrm{int}(\tilde{\Delta})) \cap (\cup_{j=1}^m \gamma_j) = \varnothing$.
Assume that $\varphi_{|\mathrm{int}(\tilde{\Delta})}$ is not an embedding, then there exists $D, E \in \mathrm{int}(\tilde{\Delta})$ such that $\varphi(D) = \varphi(E)$. Set $\overrightarrow{v} = \overrightarrow{DE} \in \mathbb{R}^2$. Since M is a translation surface, if $P, Q \in \tilde{\Delta}$ satisfy $\overrightarrow{PQ} = \pm\overrightarrow{v}$, then we have

$\varphi(P) = \varphi(Q)$. It is not difficult to see that there exists a point $P \in \tilde{\Delta} \setminus \{\tilde{A}, \tilde{B}, \tilde{P}_{j_0}\}$ such that one of the following holds

$$\begin{cases} \overrightarrow{P\tilde{P}_{j_0}} & = & \pm\overrightarrow{v}, \\ \overrightarrow{P\tilde{A}} & = & \pm\overrightarrow{v}, \\ \overrightarrow{P\tilde{B}} & = & \pm\overrightarrow{v}. \end{cases}$$

Since $\varphi(P)$ must be a regular point of the flat metric, we get a contradiction. Therefore, we can conclude that the restriction of φ to $\mathrm{int}(\tilde{\Delta})$ is an embedding.

□

Let γ' and γ'' denote the saddle connections corresponding to the sides AP_{j_0} and BP_{j_0} of Δ. By construction, $\eta^+_{j_0}$ is contained in the triangle Δ, and all the other segments in the family $\{\eta^+_j,\ j \neq j_0\}$ intersect either γ' or γ''. We can now apply the same arguments to γ' and γ'' and continue the procedure until we get a family of embedded triangles that covers all the segment η^+_j.

Remark 4.3. *Let us make the following important observation on the sides of the triangles we have constructed: if δ is a side of one of the triangles above, δ is not contained in the family $\{\gamma_1, \dots, \gamma_m\}$, then δ is one side of an embedded vertical trapezoid $\mathbf{P}$ in M, whose vertical sides corresponds to two vertical segments η^+_i, η^+_j, and the opposite side of δ in $\mathbf{P}$ is a subsegment Q_iQ_j of γ. We will call Q_iQ_j the* projection *of δ on γ. Let δ' be another side of one of the triangles we obtained, and $Q_{i'}Q_{j'}$ be its projection on γ. By construction, if Q_iQ_j and $Q_{i'}Q_{j'}$ intersect then either $Q_iQ_j \subset Q_{i'}Q_{j'}$, or $Q_{i'}Q_{j'} \subset Q_iQ_j$.*

By a symmetric procedure, we can find a family of embedded triangles, whose vertices are singularities of the flat metric, with disjoint interiors that covers all the vertical rays that intersect γ from the lower side. Note that for this case, we choose j_0 to be the largest index such that $|\eta^-_{j_0}| = \min\{|\eta^-_j|,\ j = 1, \dots, \ell\}$, where η^-_j are the lower rays at γ.

Applying this construction to all of the saddle connections in the family $\{\gamma_1, \dots, \gamma_m\}$, we get a collection of embedded triangles $\{\Delta_\alpha,\ \alpha \in I\}$ in M with vertices in the set $\{p_1, \dots, p_n\}$.

Claim 4.4. *The family $\{\Delta_\alpha, \alpha \in I\}$ is a triangulation of M.*

Proof. We first show that if Δ_{α_1} and Δ_{α_2} are two triangles in this family then $\mathrm{int}(\Delta_{\alpha_1}) \cap \mathrm{int}(\Delta_{\alpha_2}) = \varnothing$. Assume that $\mathrm{int}(\Delta_{\alpha_1}) \cap \mathrm{int}(\Delta_{\alpha_2}) \neq \varnothing$, since a triangle in $\{\Delta_\alpha, \alpha \in I\}$ cannot be included in the other one, there must exist a side δ_1 of Δ_{α_1}, and a side δ_2 of Δ_{α_2} such that $\mathrm{int}(\delta_1) \cap \mathrm{int}(\delta_2) \neq \varnothing$. By construction (see Remark 4.3) δ_i is one side of an embedded vertical trapezoid $\mathbf{P}_i$. Note that the endpoints of δ_i are singularities of M, thus they cannot be included in $\mathrm{int}(\mathbf{P}_i)$. Recall that opposite side of δ_i in $\mathbf{P}_i$ is a subsegment of a saddle connection in the family $\{\gamma_1, \dots, \gamma_m\}$. By construction, δ_i and the vertical sides of $\mathbf{P}_i$ do not intersect the set $\sqcup_{1 \leq i \leq m} \mathrm{int}(\gamma_i)$. Using these properties, one can easily check that if $\mathrm{int}(\delta_1) \cap \mathrm{int}(\delta_2) \neq \varnothing$ then we have a contradiction.

It remains to show that the union of all these triangles is M. Let M' be the closure in M of the set $M \setminus (\cup_{\alpha\in I}\Delta_\alpha)$. If $M' \neq \varnothing$, then M' is a flat surface with piecewise geodesic boundary. Let V' be the finite subset of M' which arises from the singularities of M. We can triangulate M' by geodesic segments with endpoints in V'. Consider a triangle Δ in this triangulation. We can embed isometrically Δ into $\mathbb{R}^2$ such that ξ is mapped to the vertical vector field ∂_y. Consider the vertical rays starting from the vertices of Δ, we see that there always exists a ray that intersects $\mathrm{int}(\Delta)$. This intersection corresponds to a subsegment of a separatrix in direction $\pm\xi$. But by construction, such a subsegment must be contained in the union of $\{\Delta_\alpha,\ \alpha \in I\}$, and we have a contradiction. The claim is then proven.

$\square$

We will call this triangulation constructed above the *special triangulation* of $(M, \omega, \{\gamma_1, \dots, \gamma_m\})$. By construction, this triangulation is unique. Note that such a triangulation only exists for surfaces in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$.

4.2. **Characterizing special triangulation.** Let Γ be the dual graph of the special triangulation T of $(M, \omega, \{\gamma_1, \dots, \gamma_m\})$, the vertices of Γ are the triangles in $\mathrm{T}^{(2)}$, and the (geometric) edges of Γ are the edges in $\mathrm{T}^{(1)}$. Remark that Γ is a trivalent graph, with N_2 vertices and N_1 edges. At each vertex of Γ, we have a cyclic ordering on the set of edges incident to this vertex which is induced by the orientation of the surface.

For any γ_i, the union of the triangles that covers the vertical separatrices that intersect γ_i from the upper side (resp. lower side) is dual to a subgraph of Γ which is a tree, we denote this tree by $\Gamma_{i,+}$ (resp. $\Gamma_{i,-}$), and choose its *root* to be the vertex dual to the triangle containing γ_i. By construction, all the trees $\Gamma_{i,\pm}$ are disjoint, and any vertex of Γ belongs to one of those trees. Note that if no separatrix in direction $-\xi$ (resp. ξ) reaches $\mathrm{int}(\gamma_i)$, then the tree $\Gamma_{i,+}$ (resp. $\Gamma_{i,-}$) is empty. Let us define

Definition 4.5. *Let Γ be a trivalent graph of N_2 vertices and N_1 edges, the set of edges incident to each vertex is equipped with a cyclic ordering. Let $\Gamma_{i,\varepsilon},\ i \in \{1, \dots, m\}$, $\varepsilon \in \{\pm\}$ be $2m$ disjoint subgraphs of Γ, each of which is a tree with a specified root ($\Gamma_{i,\varepsilon}$ can be empty). We will say that $(\Gamma, \{\Gamma_{i,\varepsilon}\})$ is an* admissible family of graphs *if the following conditions are satisfied.*

- *any vertex of Γ belongs to one of the trees $\Gamma_{i,\varepsilon}$,*
- *for each $i \in \{1, \dots, m\}$ the roots of $\Gamma_{i,+}$ and $\Gamma_{i,-}$ are connected by an edge of Γ.*

By definition if $(\Gamma, \{\Gamma_{i,\pm}\})$ arises from the special triangulation of $(M, \omega, \{\gamma_1, \dots, \gamma_m\})$, then $(\Gamma, \{\Gamma_{i,\pm}\})$ is an admissible family of graphs. Let us now choose a numbering of the edges of Γ as follows:

(a) we assign the edges dual to $\{\gamma_1, \dots, \gamma_m\}$ to be $\{e_1, \dots, e_m\}$ respectively,
(b) we next number the edges in the trees $\Gamma_{i,\pm}$ subsequently in such a way that , for each $\Gamma_{i,\varepsilon}$
  (b.1) if e_{i_1}, e_{i_2} belong to $\Gamma_{i,\varepsilon}$, and the distance from e_{i_1} to the root is smaller than the distance from e_{i_2} to the root then $i_1 < i_2$,



FIGURE 1. Special triangulation and dual graphs of a surface in $\widetilde{\mathcal{H}}^{(2)}(2)$.

- (b.2) if $e_{i_1}, e_{i_2}, e_{i_3}$, with $i_1 < i_2 < i_3$, belong to $\Gamma_{i,\varepsilon}$, and $e_{i_1}, e_{i_2}, e_{i_3}$ correspond to three sides of the same triangle Δ, then the cyclic order at the vertex of $\Gamma_{i,\varepsilon}$ dual to Δ is $(e_{i_1}, e_{i_2}, e_{i_3})$.

(c) We finally number the remaining edges arbitrarily.

In particular, our numbering satisfies the following conditions

(i) the first m edges are the duals of $\{\gamma_1, \dots, \gamma_m\}$,
(ii) if e_{i_1} and e_{i_2} belong to one of the trees $\Gamma_{i,\varepsilon}$, and e_{i_1} is contained in the path from e_{i_2} to the root, then $i_1 < i_2$,
(iii) if e_{i_1} belongs to one of the tree $\Gamma_{i,\varepsilon}$, and $e_{i_2} \notin \{e_1, \dots, e_m\}$ does not belong to any tree, then $i_1 < i_2$.

We will call a numbering of the edges of Γ satisfying the conditions above a *compatible numbering* with respect to the trees $\{\Gamma_{i,\pm}\}$.

By a slight abuse of notation, in what follows we will denote by the same letter an edge of Γ and its dual in $\mathrm{T}^{(1)}$. For any edge e_i in $\mathrm{T}^{(1)}$, let $z_i = x_i + \imath y_i$ be the complex number associated to e_i (by an embedding of a neighborhood of e_i in $\mathbb{R}^2$ such that ξ is mapped to ∂_y). Recall that we have $\mathrm{card}(\mathrm{T}^{(1)}) = N_1$ and $\mathrm{card}(\mathrm{T}^{(2)}) = N_2$. By assumption, we have $x_i \neq 0$, $\forall i \in \{1, \dots, N_1\}$. We always choose the orientation of e_i so that $x_i > 0$.

Set $Z = (z_1, \dots, z_{N_1}) \in \mathbb{C}^{N_1}$. Each triangle in $\mathrm{T}^{(2)}$ (vertices of Γ) gives an equation of the form (3). The collection of all of these equations gives us a system $\mathbf{S}_\mathrm{T}$. Let V_T denote the subspace of $\mathbb{C}^{N_1}$ consisting of solutions of $\mathbf{S}_\mathrm{T}$. Recall that from Lemma 3.1, we know that $\dim_\mathbb{C} \mathrm{V}_\mathrm{T} = 2g+n-1$. Clearly we have $Z \in \mathrm{V}_\mathrm{T}$. We wish now to describe the additional conditions that Z must satisfy. For this, let us pick a triangle Δ in $\mathrm{T}^{(2)}$, and let $(e_{i_1}, e_{i_2}, e_{i_3})$ be the sides of Δ written in the cyclic order induced by the orientation of M, where $i_1 = \min\{i_1, i_2, i_3\}$. Let Γ_α be the tree in the family $\{\Gamma_{i,\pm}\}$ that contains the vertex of Γ dual to Δ.

(A) If Δ is the root of Γ_α, then $e_{i_1} \in \{e_1, \dots, e_m\}$. If $e_{i_1} \notin \{e_1, \dots, e_m\}$, then e_{i_1} is the unique edge of Γ_α that is included in the path from Δ to the root. By construction, in both cases we always have

$$0 < x_{i_2} < x_{i_1} \text{ and } 0 < x_{i_3} < x_{i_1}. \tag{7}$$

We will call e_{i_1} the *base* of Δ.

(B) Given $v = (x, y)$ and $v' = (x', y')$ two vectors in $\mathbb{R}^2$, let us define

$$v \wedge v' = \det\left(\begin{smallmatrix} x & x' \\ y & y' \end{smallmatrix}\right) = xy' - x'y.$$

Considering a complex number z as a vector in $\mathbb{R}^2$, we have

$$\mathrm{A}(\Delta) = \frac{1}{2} z_{i_2} \wedge z_{i_1} = \frac{1}{2}\mathrm{Im}(z_{i_1}\overline{z}_{i_2}).$$

Thus, Z must satisfy

$$\mathrm{Im}(z_{i_1}\overline{z}_{i_2}) = x_{i_2}y_{i_1} - x_{i_1}y_{i_2} > 0. \tag{8}$$

(C) Removing e_{i_1} from the tree Γ_α, we get two subtrees Γ'_α and Γ''_α, where Γ'_α is the one that contains Δ. The union of triangles of T dual to the vertices of Γ'_α can be identified with a polygon **P** in $\mathbb{R}^2$ which has a side corresponding to e_{i_1}. Note that if $\alpha = (., +)$, then e_{i_1} is the lower side of **P** (that is all vertical downward rays from the vertices of **P** intersect e_{i_1}), and if $\alpha = (., -)$, then e_{i_1} is the upper side of **P** (all vertical upward rays from the vertices of **P** intersect e_{i_1}).

Without loss of generality, we can assume that e_{i_1} is the lower side of **P**. The triangle Δ is identified with a triangle included in **P** whose vertices are denoted by A, B, C, where A and B are the left and right endpoint of e_{i_1} respectively. Note that by the convention on the cyclic ordering, we have $e_{i_2} = CB, e_{i_3} = AC$.

If P is a vertex of **P** different from A and B, we denote by $\hat{P}$ the intersection of the vertical (downward) ray from P with $\mathrm{int}(AB)$, we will call $\hat{P}$ the projection of P to AB. Let us denote by $h(P)$ the length of the segment $P\hat{P}$. We denote the vertices of **P** whose projection to AB is between A and $\hat{C}$ by $D_1, \dots, D_r$ so that $\hat{D}_i$ is between A and $\hat{D}_{i+1}$. Similarly, we denote the vertices of **P** whose projection to AB is between $\hat{C}$ and B by $E_1, \dots, E_s$ in such a way that $\hat{E}_j$ is between $\hat{C}$ and $\hat{E}_{j+1}$.

Recall that by construction, we have

$$h(C) < h(D_i), \ \forall i = 1, \dots, r,$$

and

$$h(C) \leq h(E_j), \ j = 1, \dots, s.$$

It is easy to check that the condition $h(C) < h(D_i)$ is equivalent to $\overrightarrow{D_iC} \wedge \overrightarrow{AB} > 0$. As $\overrightarrow{D_iC} = \overrightarrow{D_iD_{i+1}} + \cdots + \overrightarrow{D_rC}$, and the sides of **P** are the edges of T, we can write $\overrightarrow{D_iC}$ as a linear function f_i of Z. Consequently, the condition on $h(D_i)$ becomes

$$\mathrm{Im}(z_{i_1}\overline{f_i}) > 0, \ \forall i = 1, \dots, r. \tag{9}$$

Similarly, for $E_j, \ j = 1, \dots, s$, we have $h(C) \leq h(E_j)$, and these conditions are equivalent two

$$\mathrm{Im}(z_{i_1}\overline{g_j}) \geq 0, \ \forall j = 1, \dots, s. \tag{10}$$

where g_j are some linear functions of Z. The functions f_i, g_j are completely determined by the tree Γ'_α. Note also that, *a priori*, we cannot replace the inequality in (10) by a strict one.

Let $(\Gamma, \{\Gamma_{i,\pm}\})$ be an admissible family of graphs, and choose a compatible numbering of the edges of Γ. Let V be the space of solutions in $\mathbb{C}^{N_1}$ of the system of linear equations associated to the vertices of Γ. Let $\mathcal{D}$ be the subset of V which is defined by the inequations (7), (8), (9), (10). For every $Z = (z_1, \dots, z_{N_1}) \in \mathcal{D}$, we define $\Psi(Z)$ to be the triple $(M, \omega, \{\gamma_1, \dots, \gamma_m\})$, where

- M is the surface obtained by gluing the triangles constructed from the coordinates of Z (using the graph Γ).
- ω is the holomorphic one-form on M satisfying $\omega = dz$ in each triangle.
- γ_i is the saddle connection in Σ corresponding to the edge of Γ which connects the roots of $\Gamma_{i,+}$, and $\Gamma_{i,-}$.

Proposition 4.6. *Let $\mathcal{D}_0$ be a component of $\mathcal{D}$. If $\Psi(Z) \in \widetilde{\mathcal{H}}^{(m)}(\underline{k})$ for some $Z \in \mathcal{D}_0$, then $\Psi(\mathcal{D}_0) \subset \widetilde{\mathcal{H}}^{(m)}(\underline{k})$, and Ψ realizes a homeomorphism from $\mathcal{D}_0$ onto its image.*

Proof. Since the topological invariants of $\Psi(Z)$ depends continuously on Z, they are constant on each connected component of $\mathcal{D}$.

If $(M, \omega, \{\gamma_1, \dots, \gamma_m\}) = \Psi(Z)$, the surface M can be constructed as follows: for each tree $\Gamma_{i,\varepsilon}$, we construct the polygon $\mathbf{P}_{i,\varepsilon} \in \mathbb{R}^2$ by gluing the triangles corresponding to the vertices of $\Gamma_{i,\varepsilon}$, we then glue the polygons $\mathbf{P}_{i,\varepsilon}$ to get M. Note that by construction, we obtain M together with an admissible triangulation T whose dual graph is Γ. Since Z satisfies the inequations (7), (8), (9), (10), T is the special triangulation of $(M, \omega, \{\gamma_1, \dots, g_m\})$. If we have $\Psi(Z) = \Psi(Z')$ but $Z' \neq Z$, then we obtain an element in $\widetilde{\mathcal{H}}^{(m)}(\underline{k})$ with two special triangulations, which is impossible. It is clear that Ψ is a locally homeomorphic. The proposition is then proven. □

Since $\mathcal{D}$ is defined by finitely many inequations, it has only finitely many components. Remark that the action of the subgroup $U_- = \left(\begin{smallmatrix} * & 0 \\ * & * \end{smallmatrix}\right)$ of $\mathrm{GL}^+(2, \mathbb{R})$ on $\mathbb{C}^{N_1}$ preserves the inequations (7), (8), (9), (10). Therefore, we have shown the following

Theorem 4.7. *There exists a partition of* $\widetilde{\mathcal{H}}^{(m)}(\underline{k})^*$ *into finitely many subsets which are* U_-*-invariant. Each subset in this partition is homeomorphic to a component of the subset of a linear subspace of dimension* $2g+n-1$ *of* $\mathbb{C}^{N_1}$*, which is determined by the inequations (7), (8), (9), (10) via the map* Ψ*.*

5. PROOF OF THEOREM 2.3

5.1. **Primary and auxiliary families of indices.** Given an admissible family of graphs $(\Gamma, \{\Gamma_{i,\pm}\})$, where Γ is the dual graph of a special triangulation T of an element $(M, \omega, \{\gamma_1, \dots, \gamma_m\}) \in \widetilde{\mathcal{H}}^{(m)}(\underline{k})$, let us choose a compatible numbering of the edges of Γ. We denote by $\mathbf{S}_\Gamma$ the systems of linear equations of type (3) associated to Γ, and by V_Γ the subspace of $\mathbb{C}^{N_1}$ consisting of solution of $\mathbf{S}_\Gamma$.

In what follows, we will say that a family of coordinates $\{z_i, i \in I\}$ is *dependent* with respect to $\mathbf{S}_\Gamma$ if there exists a vector $(\lambda_i)_{i\in I} \in \mathbb{R}^I$ such that $\sum_{i\in I} \lambda_i z_i = 0$, for all $(z_1, \dots, z_{N_1}) \in \mathrm{V}_\Gamma$. The family $\{z_i,\ i \in I\}$ is said to be *independent* with respect to $\mathbf{S}_\Gamma$, if such a vector does not exist.
We have another interpretation of dependent and independent families as follows: let $a_i, i = 1, \dots, N_2$, denote the row vectors corresponding to the equations of the system $\mathbf{S}_\Gamma$, we consider a_i as an element of the space $(\mathbb{C}^{N_1})^*$ which is equivariant by conjugation. Set $\mathrm{V}^*_\Gamma = \mathrm{Vect}_\mathbb{R}\{a_1, \dots, a_{N_2}\} \subset (\mathbb{C}^{N_1})^*$. For any family $I \subset \{1, \dots, N_1\}$, set $\mathbb{R}^I = \{(\lambda_1, \dots, \lambda_{N_1}) : \lambda_i = 0 \text{ if } i \notin I\} \subset \mathbb{R}^{N_1}$, we regard $\mathbb{R}^I$ as a subspace of the space of $\mathbb{C}$-linear forms $(\mathbb{C}^{N_1})^*$ equivariant by conjugation. If $a = (\lambda_i)_{i\in I} \in \mathbb{R}^I \cap \mathrm{V}^*_\Gamma$, then $a(Z) = \sum_{i\in I} \lambda_i z_i = 0$ for all $Z = (z_1, \dots, z_{N_1}) \in \mathrm{V}_\Gamma$. Thus I is an independent family if and only if $\mathbb{R}^I \cap \mathrm{V}^*_\Gamma = \{0\}$.

Remark 5.1.

- *Since the coefficients of* a_i *are in* $\{0, \pm 1\}$*, if the family* $\{z_i,\ i \in I\}$ *is dependent, we can choose the vector* $(\lambda_i)_{i\in I}$ *such that* $\lambda_i \in \mathbb{Z}$*, for all* $i \in I$*.*
- *Let* e_i *denote the edge of* T *which corresponds to* z_i*. A family* I *is independent with respect to* $\mathbf{S}_\Gamma$ *if and only if the family* $\{e_i,\ i \in I\}$ *is independent in* $H_1(M, \{p_1, \dots, p_n\}; \mathbb{Z})$*.*

Definition 5.2. *The* primary family *of indices of* $(\Gamma, \{\Gamma_{i,\pm}\})$ *is the unique ordered subset* $(i_1, \dots, i_{2g+n-1})$ *of* $\{1, \dots, N_1\}$ *that satisfies the following properties*

- $i_k < i_{k+1}$*,*
- $i_1, \dots, i_{2g+n-1}$ *is an independent family of indices with respect to* $\mathbf{S}_\Gamma$*,*
- *For all* $k \in \{1, \dots, 2g+n-1\}$*, if* $i < i_k$ *then the family* $(i_1, \dots, i_{k-1}, i)$ *is dependent with respect to* $\mathbf{S}_\Gamma$*.*

Remark 5.3. *The primary family of indices can be found inductively by the following procedure: first we take* $i_1 = 1, \dots, i_m = m$*, recall that* $e_1 = \gamma_1, \dots, e_m = \gamma_m$*, and by assumption* $(\gamma_1, \dots, \gamma_m)$ *is independent in* $H_1(M, \{p_1, \dots, p_n\}; \mathbb{Z})$*, hence the family* $\{i_1, \dots, i_m\}$ *is independent with respect to* $\mathbf{S}_\Gamma$*. Assume that we already have an independent family* $(i_1, \dots, i_k)$*, then* i_{k+1} *is the smallest index such that* $(i_1, \dots, i_k, i_{k+1})$ *is independent.*

Let us denote by I the primary family of index of $(\Gamma, \{\Gamma_{i,\pm}\})$.

Definition 5.4. *An* auxiliary family *of I is an ordered family of indices $J = (j_{m+1}, \dots, j_{2g+n-1})$ which satisfies the following condition: for any $k \in \{m+1, \dots, 2g+n-1\}$, and e_{j_k} is the base of one of the triangles in* T *that contains e_{i_k}. In particular, we have $j_k < i_k$.*

Remark 5.5. *By definition, we have $j_k < i_k$, thus $(i_1, \dots, i_{k-1}, j_k)$ is a dependent family. Since the family $(i_1, \dots, i_{k-1})$ is independent, it follows that z_{j_k} is a linear function of $(z_{i_1}, \dots, z_{i_{k-1}})$ in V_Γ.*

An auxiliary family of I can be found as follows: for each $i_k \in I$, $k > m$, consider the corresponding edge e_{i_k} of Γ. We have two cases:

- e_{i_k} belongs to some tree Γ_α. In this case, let Δ_k be the triangle of T dual to the endpoint of e_{i_k} which is closer to the root of Γ_α. Let $e_{i'_k}$ and $e_{i''_k}$ be the other sides of Δ_k where $i'_k > i''_k$. We have $i_k \neq \min\{i_k, i'_k, i''_k\}$ by the definition of compatible numbering. Thus $i''_k = \min\{i_k, i'_k, i''_k\}$, which implies that $e_{i''_k}$ is the base of Δ_k, and we can choose j_k to be i''_k.
- e_{i_k} does not belong to any of the trees Γ_α. Let Δ_k be one of the two triangles in T that contains e_{i_k}. Again let $e_{i'_k}$ and $e_{i''_k}$ be the other sides of Δ_k where $i'_k > i''_k$. By the definition of compatible numbering, we also have $i_k \neq \min\{i_k, i'_k, i''_k\}$, hence we can choose j_k to be i''_k.

5.2. **Proof of Theorem 2.3.**

Proof. Fix an admissible family of graphs $(\Gamma, \{\Gamma_{i,\pm}\})$ together with a compatible numbering of the edges of Γ. Let $I = (i_1, \dots, i_{2g+n-1})$ be the primary family of indices of $(\Gamma, \{\Gamma_{i,\pm}\})$ and $J = (j_{m+1}, \dots, j_{2g+n-1})$ an auxiliary family of indices of I. Since $\mathrm{card}\{I\} = \dim_{\mathbb{C}} \mathrm{V}_\Gamma$, there exists a linear isomorphism with rational coefficients

$$\begin{array}{cccc} \mathrm{F}: & \mathbb{C}^{2g+n-1} & \to & \mathrm{V}_\Gamma \\ & (z_1, \dots, z_{2g+n-1}) & \mapsto & (f_1(z_1, \dots, z_{2g+n-1})), \dots, f_{N_1}(z_1, \dots, z_{2g+n-1})) \end{array}$$

which satisfies $f_{i_k} = z_k$, $\forall i_k \in I$. To simplify the notations, we write f_k instead of f_{j_k}, for every $j_k \in J$.

Let $\mathcal{D}$ be the domain of V_Γ which is defined by the inequations (7), (8), (9), (10). By a slight abuse of notations, we will denote also by $\mathcal{D}$ the corresponding domain in $\mathbb{C}^{2g+n-1}$ (via F). For every $(z_1, \dots, z_{2g+n-1}) \in \mathcal{D}$, let $\mathrm{A}((z_1, \dots, z_{2g+n-1}))$ denote the area of the surface constructed from $F((z_1, \dots, z_{2g+n-1}))$.

Let us write $z_i = x_i + \imath y_i$, with $x_i, y_i \in \mathbb{R}$, $i = 1, \dots, 2g+n-1$, and $f_k = a_k + \imath b_k$, $a_k, b_k \in \mathbb{R}$, $k = m+1, \dots, 2g+n-1$. Remark that the coefficients of f_k are real, therefore a_k is a real linear function of $(x_1, \dots, x_{k-1})$, and b_k is a real linear function of $(y_1, \dots, y_{k-1})$.

To prove the theorem, is is enough to show that

$$\int_{\mathcal{D}} e^{-(|z_1|^2/\epsilon_1^2 + \dots + |z_m|^2/\epsilon_m^2)} e^{-\mathrm{A}} dx_1 dy_1 \dots dx_{2g+n-1} dy_{2g+n-1} < K\epsilon_1^2 \dots \epsilon_m^2. \tag{11}$$

To prove the inequality (11) we will only make use of the conditions (7), and (8). We first observe that the inequation (7) implies that for any $(x_1 + \imath y_1, \dots, x_{2g+n-1} + \imath y_{2g+n-1}) \in \mathcal{D}$, we have

$$0 < x_k < a_k, \ k = m+1, \dots, 2g+n-1. \tag{12}$$

Set

$$\mathcal{D}^* = \{(\underline{z}, \underline{x}), \ \underline{z} \in \mathbb{C}^m, \underline{x} \in \mathbb{R}_+^{2g+n-1-m}, \ (\underline{z}, \underline{x}) \text{ satisfies (12)}\} \subset \mathbb{C}^m \times \mathbb{R}_+^{2g+n-1-m}.$$

For any $(\underline{z}, \underline{x}) \in \mathcal{D}^*$, set

$$\mathcal{Y}(\underline{z}, \underline{x}) = \{\underline{y} = (y_{m+1}, \dots, y_{2g+n-1}) \in \mathbb{R}^{2g+n-1-m} : \ (\underline{z}, \underline{x} + \imath\underline{y}) \in \mathcal{D}\}.$$

The inequality (11) now becomes

$$\mathcal{I} := \int_{\mathcal{D}^*} e^{-(|z_1|^2/\epsilon_1^2+\cdots+|z_m|^2/\epsilon_m^2)} \left(\left(\int_{\mathcal{Y}(\underline{z},\underline{x})} e^{-\mathrm{A}} dy_{m+1} \dots dy_{2g+n-1} \right) d\lambda_{2g+n-1-m} \right) d\lambda_{2m} < K\epsilon_1^2 \dots \epsilon_m^2. \tag{13}$$

where $\lambda_{2g+n-1-m} = dx_{m+1} \dots dx_{2g+n-1}$ and $d\lambda_{2m} = dx_1 dy_1 \dots dx_m dy_m$ are the Lebesgue measures of $\mathbb{R}^{2g+n-1-m}$ and $\mathbb{C}^m$ respectively.

For every $k \in \{m+1, \dots, 2g+n-1\}$, let Δ_k denote the triangle in T that contains e_{i_k} and e_{j_k}.

Claim 5.6. *If $k \neq k'$, then $\Delta_k \neq \Delta_{k'}$.*

Proof. We can assume that $k < k'$. If $\Delta_k = \Delta_{k'}$ then $e_{i_k}, e_{j_k}, e_{i_{k'}}$ are contained in the same triangle, which implies that $z_{i_{k'}}$ is a linear function of (z_{i_k}, z_{j_k}) in V_Γ. By assumption, z_{j_k} is a linear function of $(z_{i_1}, \dots, z_{i_{k-1}})$, thus $z_{i_{k'}}$ is a linear function of $(z_{i_1}, \dots, z_{i_k})$ which is impossible since $(i_1, \dots, i_{k'})$ is an independent family. □

Let η_k be the area of Δ_k. The previous claim implies immediately that

$$A > \sum_{k=m+1}^{2g+n-1} \eta_k. \tag{14}$$

As functions on $\mathbb{C}^{2g+n-1}$, η_k are given by

$$\eta_k = \pm\frac{1}{2}\mathrm{Im}(f_k\bar{z}_k) = \pm\frac{1}{2}(a_k y_k - x_k b_k).$$

Since b_k is a linear function of $(y_1, \dots, y_{k-1})$, we have

$$\frac{\partial\eta_k}{\partial y_j} = \begin{cases} \frac{\pm1}{2} x_k \frac{\partial b_k}{\partial y_j} & \text{if } j < k, \\ \frac{\pm1}{2} a_k & \text{if } j = k, \\ 0 & \text{if } j > k, \end{cases}$$

and

$$d\eta_{m+1}\dots d\eta_{2g+n-1} = \frac{1}{2^{2g+n-1-m}}|\det\begin{pmatrix} \pm a_{m+1} & 0 & \dots & 0 \\ * & \pm a_{m+2} & \dots & 0 \\ \dots & \dots & \dots & \dots \\ * & * & \dots & \pm a_{2g+n-1} \end{pmatrix}|dy_{m+1}\dots dy_{2g+n-1}.$$

Therefore, we have

$$dy_{m+1}\dots dy_{2g+n-1} = \frac{2^{2g+n-1-m}}{a_{m+1}\dots a_{2g+n-1}} d\eta_{m+1}\dots d\eta_{2g+n-1}.$$

It follows from (14) that

$$\mathcal{I}(\underline{z},\underline{x}) := \int_{\mathcal{Y}(\underline{z},\underline{x})} e^{-A} dy_{m+1}\dots dy_{2g+n-1} < \frac{2^{2g+n-1-m}}{a_{m+1}\dots a_{2g+n-1}} \int_{\mathcal{Y}(\underline{z},\underline{x})} e^{-(\eta_{m+1}+\dots+\eta_{2g+n-1})} d\eta_{m+1}\dots d\eta_{2g+n-1}.$$

The conditions (8) mean that the functions η_k are positive on $\mathcal{Y}(\underline{z},\underline{x})$. Thus we have

$$\mathcal{I}(\underline{z},\underline{x}) < \frac{2^{2g+n-1-m}}{a_{m+1}\dots a_{2g+n-1}} \int_0^{+\infty} e^{-\eta_{m+1}} d\eta_{m+1}\dots \int_0^{+\infty} e^{-\eta_{2g+n-1}} d\eta_{2g+n-1} = \frac{2^{2g+n-1-m}}{a_{m+1}\dots a_{2g+n-1}},$$

and

$$\mathcal{I} < 2^{2g+n-1-m} \int_{\mathcal{D}^*} \frac{e^{-(|z_1|^2/\epsilon_1^2+\dots+|z_m|^2/\epsilon_m^2)}}{a_{m+1}\dots a_{2g+n-1}} dx_{2g+n-1}\dots dx_{m+1} d\lambda_{2m}. \tag{15}$$

From the definition of $\mathcal{D}^*$, the right hand side of (15) is equal to

$$2^{2g+n-1-m} \int_{\mathbb{C}^m} e^{-(|z_1|^2/\epsilon_1^2+\dots+|z_m|^2/\epsilon_m^2)} \left(\int_0^{a_{m+1}} \left(\dots\left(\int_0^{a_{2g+m-1}} \frac{1}{a_{m+1}\dots a_{2g+n-1}} dx_{2g+n-1}\right)\dots\right) dx_{m+1}\right) d\lambda_{2m}.$$

Since $a_{m+1},\dots,a_k$ do not depend on x_k, we have

$$\int_0^{a_{m+1}} \left(\dots\left(\int_0^{a_{2g+m-1}} \frac{1}{a_{m+1}\dots a_{2g+n-1}} dx_{2g+n-1}\right)\dots\right) dx_{m+1} = 1.$$

Thus

$$\mathcal{I} < 2^{2g+n-1-m} \int_{\mathbb{C}^m} e^{-(|z_1|^2/\epsilon_1^2+\dots+|z_m|^2/\epsilon_m^2)} dx_1 dy_1 \dots dx_m dy_m = 2^{2g+n-1-m} \pi^m \epsilon_1^2 \dots \epsilon_m^2.$$

The inequality (13) is then proven. The proof of Theorem 2.3 is now complete.

□

6. QUADRATIC DIFFERENTIALS

6.1. **Half-translation surface with marked saddle connections.** Let us denote by $\widetilde{Q}^{(m)}(\underline{d})$ the set of triple $(M,\phi,\{\gamma_1,\dots,\gamma_m\})$, where

- (M,ϕ) is an element of $Q(\underline{d})$,
- $\{\gamma_1,\dots,\gamma_m\}$ be an independent family of saddle connections in M.

There exists a natural projection $\varrho : \widetilde{Q}^{(m)}(\underline{d}) \to Q(\underline{d})$ consisting of forgetting the markings of the saddle connections. This projection is locally homeomorphic, hence we can pullback the Lebesgue measure $d\mu$ of $Q(\underline{d})$ to $\widetilde{Q}^{(m)}(\underline{d})$. We define a function $\mathfrak{F}_{\underline{\epsilon}} : \widetilde{Q}^{(m)}(\underline{d}) \to \mathbb{R}$ as follows

$$\mathfrak{F}_{\underline{\epsilon}} : (M,\phi,\{\gamma_1,\dots,\gamma_m\}) \mapsto \exp(-\sum_{j=1}^{m}\frac{|\gamma_j|}{\epsilon_j^2} - \mathrm{A}(M)).$$

Theorem 1.4 is a direct consequence of the following theorem

Theorem 6.1. *There exists a constant $K = K(\underline{d},m)$ such that*

$$\int_{\widetilde{Q}^{(m)}(\underline{d})} \mathfrak{F}_{\underline{\epsilon}}d\mu < K\epsilon_1^2\dots\epsilon_m^2.$$

6.2. **Proof of Theorem 1.4.**

Proof. Let $Q^{(m)}_{\underline{\epsilon}}(\underline{d})$ be the subset of $Q(\underline{d})$ consisting of pairs (M,ϕ) such that there exists an independent family of m saddle connections $\{\gamma_1,\dots,\gamma_m\}$ on M satisfying $|\gamma_j|^2 < \epsilon_j^2\mathrm{A}(M)$. Let $\widetilde{Q}^{(m)}_{\underline{\epsilon}}(\underline{d})$ be the pre-image of $Q^{(m)}_{\underline{\epsilon}}(\underline{d})$ by the natural projection $\varrho : \widetilde{Q}^{(m)}(\underline{d}) \to Q(\underline{d})$. Theorem 6.1 implies immediately that

$$\int_{\widetilde{Q}^{(m)}_{\underline{\epsilon}}(\underline{d})} \mathfrak{F}_{\underline{\epsilon}}d\mu < K\epsilon_1^2\dots\epsilon_m^2. \tag{16}$$

Set

$$\begin{array}{rccl} f_{\underline{\epsilon}}: & Q(\underline{d}) & \to & \mathbb{R} \\ & (M,\phi) & \mapsto & e^{-\mathrm{A}(M)}\left(\sum\limits_{\substack{\{\gamma_1,\dots,\gamma_m\} \\ \text{independent family in } M}} e^{-(\frac{|\gamma_1|^2}{\epsilon_1^2}+\dots+\frac{|\gamma_m|^2}{\epsilon_m^2})}\right). \end{array}$$

We have

$$\int_{\widetilde{Q}^{(m)}_{\underline{\epsilon}}(\underline{d})} \mathfrak{F}_{\underline{\epsilon}}d\mu = \int_{Q^{(m)}_{\underline{\epsilon}}(\underline{d})} f_{\underline{\epsilon}}d\mu.$$

But by definition we have $f_{\underline{\epsilon}}(.) > e^{-(m+1)\mathrm{A}(.)}$ on $Q^{(m)}_{\underline{\epsilon}}(\underline{d})$. Therefore

$$\int_{\mathcal{Q}^{(m)}_{\underline{\epsilon}}(\underline{d})} e^{-(m+1)\mathrm{A}(.)}d\mu < K\epsilon_1^2\dots\epsilon_m^2. \tag{17}$$

Now, the space $\mathcal{Q}(\underline{d})$ can be locally identified with an open in $\mathbb{R}^{2d}$, where $d = \dim_{\mathbb{C}}\mathcal{Q}(\underline{d})$, so that $d\mu$ is the Lebesgue measure. In this identification, $\mathcal{Q}_1(\underline{d})$ becomes the hypersurface $\mathbf{Q}_1$ defined by the equation $\mathrm{A} = 1$. Using the change of variables

$$\begin{array}{rccc} \Phi : & \mathbf{Q}_1\times]0,+\infty[& \to & \mathbb{R}^{2d} \\ & (v,t) & \mapsto & t\cdot v, \end{array}$$

we have $\Phi^*A = t^2$, and $\Phi^*d\mu = 2t^{2d-1}dtd\mu_1$. The inequality (17) then implies

$$\int_{\mathcal{Q}^{(m)}_{1,\underline{\epsilon}}(\underline{d})} (\int_0^{+\infty} 2t^{2d-1}e^{-(m+1)t^2}dt)d\mu_1 < K\epsilon_1^2\dots\epsilon_m^2 \tag{18}$$

Thus,

$$\mu_1(\mathcal{Q}^{(m)}_{1,\underline{\epsilon}}(\underline{d})) < C\epsilon_1^2\dots\epsilon_m^2.$$

$\square$

6.3. **Symmetric special triangulation.** Given an element $(M,\phi,\{\gamma_1,\dots,\gamma_m\})$ of $\widetilde{\mathcal{Q}}^{(m)}(\underline{d})$, let $\pi : \hat{M}\to M$ be the canonical double covering of M, and τ be the covering involution of $\hat{M}$. Let $\hat{\omega}$ be the holomorphic one-form on $\hat{M}$ such that $\pi^*\phi = \hat{\omega}^2$. We first notice that $(\hat{M},\hat{\omega})$ belongs to some stratum $\mathcal{H}(\hat{\underline{k}})$, where the vector $\hat{\underline{k}}$ depends only on the component of $\mathcal{Q}(\underline{d})$ that contains (M,ϕ). We denote by $\hat{\gamma}_i^1,\hat{\gamma}_i^2$ the pre-images of γ_i by π, which are saddle connections in $\hat{M}$. Note that the orientation of $\hat{\gamma}_i^s$ is chosen so that $\tau(\hat{\gamma}_i^1) = -\hat{\gamma}_i^2$ (which implies $\tau(\hat{\gamma}_i^2) = -\hat{\gamma}_i^1$). By assumption, we have $\mathrm{int}(\hat{\gamma}_i^s)\cap\mathrm{int}(\hat{\gamma}_{i'}^{s'}) = \varnothing$ unless $(i,s) = (i',s')$.

Given $v\in\mathbb{S}^1$, the direction v is not well defined on M, but the notion of "*parallel*" is well defined. In what follows, we will call a geodesic segment (resp. separatrix, saddle connection) parallel to the vectors $(0,\pm 1)$ a *vertical* segment (resp. separatrix, saddle connection).

Let us recall briefly a way to define the local chart on a neighborhood of $(M,\phi,\{\gamma_1,\dots,\gamma_m\})$. Set $\hat{S} = \pi^{-1}(S)$, where S is the set of zeros and poles of ϕ. Let $H_1(\hat{M},\hat{S};\mathbb{Z})^-$ be the set of cycles $c\in H_1(\hat{M},\hat{S};\mathbb{Z})$ such that $\tau(c) = -c$. A surface (M',ϕ') close to (M,ϕ) gives rise to a surface $(\hat{M}',\hat{\omega}')$ close to $(\hat{M},\hat{\omega})$ together with an involution τ' in the same homotopy class as τ. Thus we can identify $H_1(\hat{M}',\hat{S}';\mathbb{Z})^-$ with $H_1(\hat{M},\hat{S};\mathbb{Z})^-$. Choose a basis $\{c_1,\dots,c_d\}$ of $H_1(\hat{M},\hat{S};\mathbb{Z})^-$, then the period mapping $\hat{\Phi} : (M',\phi')\mapsto(\int_{c_1}\hat{\omega}',\dots,\int_{c_d}\hat{\omega}')$ is a local chart for $\mathcal{Q}(\underline{d})$, in which the volume form μ is identified with the Lebesgue measure of $\mathbb{C}^d$. If (M',ϕ') is close enough to (M,ϕ) in $\mathcal{Q}(\underline{d})$, then there exists a homeomorphism $\varphi : M\to M'$ such that $\{\varphi(\gamma_1),\dots,\varphi(\gamma_m)\}$ is a family of saddle connections with disjoint interior. Therefore, $\hat{\Phi}$ can also be viewed as a local chart for $\widetilde{\mathcal{Q}}^{(m)}(\underline{d})$ in a neighborhood of $(M,\phi,\{\gamma_1,\dots,\gamma_m\})$.

Let $\widetilde{Q}^{(m)}(\underline{d})^*$ be the set of $(M,\phi,\{\gamma_1,\dots,\gamma_m\}) \in \widetilde{Q}^{(m)}(\underline{d})$ which satisfy the following condition: every vertical separatrix intersects the set $\cup_{1\leq i\leq m}\mathrm{int}(\gamma_i)$ before reaching a singular point. The following lemma follows from the same argument as Lemma 4.1

Lemma 6.2. *$\widetilde{Q}^{(m)}(\underline{d})^*$ is an open dense subset, hence of full measure, of $\widetilde{Q}^{(m)}(\underline{d})$.*

Suppose that $(M,\phi,\{\gamma_1,\dots,\gamma_m\})$ belongs to $\widetilde{Q}^{(m)}(\underline{d})^*$, then $(\hat{M},\hat{\omega},\{\hat{\gamma}^1_1,\hat{\gamma}^2_1,\dots,\hat{\gamma}^1_m,\hat{\gamma}^2_m\})$ belongs to $\widetilde{\mathcal{H}}^{(2m)}(\underline{\hat{k}})^*$. Let $\hat{\mathrm{T}}$ be the special triangulation of $\hat{M}$ with respect to the family $\{\hat{\gamma}^s_i\}$. Since we have $\tau^*\hat{\omega} = -\hat{\omega}$, it follows that the involution τ sends a separatrix indirection $(0,1)$ to a separatrix in direction $(0,-1)$ and vice versa. By definition, the family of saddle connections $\{\hat{\gamma}^s_i\}$ is invariant under τ, therefore, τ induces a permutation on set of vertical separatrices joining singular points of $\hat{M}$ to the segments $\hat{\gamma}^s_i$. As a consequence, the triangulation $\hat{\mathrm{T}}$ is invariant under τ. We deduce in particular that τ induces an involution on the set $\hat{\mathrm{T}}^{(1)}$.

Let $\hat{N}_1$ and $\hat{N}_2$ be the number of edges and of triangles in $\hat{\mathrm{T}}$ respectively. We choose the orientations of the edges of $\hat{\mathrm{T}}$ so that $\hat{\omega}(\tau(e)) = -\hat{\omega}(e)$ for all $e \in \hat{\mathrm{T}}^{(1)}$. We consider a vector $Z \in \mathbb{C}^{\hat{N}_1}$ as a function from $\hat{\mathrm{T}}^{(1)}$ to $\mathbb{C}$. We have a system $\mathbf{S}_{\hat{\mathrm{T}}}$ of $\hat{N}_2$ linear equations of the form (3), each of which corresponding to a triangle in $\hat{\mathrm{T}}^{(2)}$. We add to this system the equations

$$Z(e') = Z(e) \tag{19}$$

where $e,e' \in \hat{\mathrm{T}}^{(1)}$ such that $\tau(e) = -e'$. Let $\hat{\mathbf{S}}_{\hat{\mathrm{T}}}$ denote the resulting system, and $\hat{\mathrm{V}}_{\hat{\mathrm{T}}} \subset \mathbb{C}^{\hat{N}_1}$ denote the space of solutions of $\hat{\mathbf{S}}_{\hat{\mathrm{T}}}$. Clearly, the integrals of $\hat{\omega}$ along the edges of $\hat{\mathrm{T}}$ gives us a vector in $\hat{\mathrm{V}}_{\hat{\mathrm{T}}}$. Given $Z' \in \hat{\mathrm{V}}_{\hat{\mathrm{T}}}$ close to Z, we can construct a surface $(\hat{M}',\hat{\omega}')$ together with $2m$ saddle connections $\hat{\gamma}'^s_i$, and an involution $\tau' : \hat{M}' \to \hat{M}'$ satisfying $\tau'(\hat{\gamma}'^1_i) = -\hat{\gamma}'^2_i$. Thus we have a continuous mapping $\Psi_{\hat{\mathrm{T}}}$ defined in a neighborhood of Z in $\hat{\mathrm{V}}_{\hat{\mathrm{T}}}$ to $\widetilde{Q}^{(m)}(\underline{d})$. It is not difficult to check that $\dim_{\mathbb{C}} \hat{\mathrm{V}}_{\hat{\mathrm{T}}} = \dim_{\mathbb{C}} \widetilde{Q}^{(m)}(\underline{d}) = 2g+n-2$, and in a local chart of $\widetilde{Q}^{(m)}(\underline{d})$, $\Psi_{\hat{\mathrm{T}}}$ is a linear isomorphism between complex vector spaces. Let $\mathcal{D}$ be the domain in $\hat{\mathrm{V}}_{\hat{\mathrm{T}}}$ which is defined by the inequations (7),(8), (9), (10), and $\mathcal{D}_0$ be the component of $\mathcal{D}$ that contains Z. The following proposition follows from the same argument as Proposition 4.6

Proposition 6.3. *The map $\Psi_{\hat{\mathrm{T}}}$ is well defined and injective in $\mathcal{D}_0$, it realizes a homeomorphism between $\mathcal{D}_0$ and its image in $\widetilde{Q}^{(m)}(\underline{d})$.*

Let us now define

Definition 6.4. *Let Γ be a trivalent graph with $\hat{N}_2$ vertices and $\hat{N}_1$ edges. Let $\Gamma^s_{i,\pm}$, $i \in \{1,\dots,m\}$, $s \in \{1,2\}$, be a family of disjoint subgraphs of Γ which are trees. We will say that $(\Gamma,\{\Gamma^s_{i,\pm}\})$ is a* symmetric admissible family of graphs *if $(\Gamma,\{\Gamma^s_{i,\pm}\})$ is an admissible family of graphs (see Definition 4.5), and there exists an involution τ of Γ which satisfies $\tau(\Gamma^s_{i,\varepsilon}) = \Gamma^{s+1}_{i,-\varepsilon}$, where we use the conventions $3 \simeq 1$, " $- -$" $\simeq$ " $+$ ", " $- +$" $\simeq$ " $-$ ".*

Let Γ be the dual graph of $\hat{\mathrm{T}}$. For each $\hat{\gamma}_i^s$, let $\Gamma_{i,+}^s$ (resp. $\Gamma_{i,-}^s$) be the tree in Γ which is dual to the union of triangles in $\hat{\mathrm{T}}^{(2)}$ that cover the vertical separatrices reaching $\mathrm{int}(\hat{\gamma}_i^s)$ from the upper side (resp. lower side). By construction, $(\Gamma, \{\Gamma_{i,\pm}^s\})$ is a symmetric admissible family of graphs. The following theorem follows from the same arguments as Theorem 4.7

Theorem 6.5. *There exists a partition of* $\widetilde{\mathcal{Q}}^{(m)}(\underline{d})^*$ *into finitely many subset, each of which corresponds to a component of the subset determined by the inequations (7),(8), (9), (10) of a subspace* $\hat{\mathrm{V}}$ *of dimension* $2g+n-2$ *of* $\mathbb{C}^{\hat{N}_1}$. *The space* $\hat{\mathrm{V}}$ *itself is determined by a symmetric admissible family of graphs.*

6.4. **Proof of Theorem 6.1.**

Proof. Let $(\Gamma, \{\Gamma_{i,\pm}^s\})$ be a symmetric admissible family of graphs. Choose a compatible numbering of the edges of Γ. Without loss of generality, we can assume that e_i is the root of $\Gamma_{i,+}^1$, $i = 1, \dots, m$. Let $\mathbf{S}_\Gamma$ denote the linear system associated to $(\Gamma, \{\Gamma_{i,\pm}^s\})$, and $\hat{\mathbf{S}}_\Gamma$ the system obtained by adding to $\mathbf{S}_\Gamma$ the equations of type (19). Let $\hat{\mathrm{V}}$ be the space of solutions of $\hat{\mathbf{S}}_\Gamma$ in $\mathbb{C}^{\hat{N}_1}$. Let $\mathcal{D}$ be the open subset of $\hat{\mathrm{V}}$ determined by the inequations (7), (8), (9), (10). We consider a vector $Z = (z_1, \dots, z_{\hat{N}_1}) \in \mathbb{C}^{\hat{N}_1}$ as a function from the edges of Γ to $\mathbb{C}$, where $z_i = Z(e_i)$. By construction, there exists a continuous map $\Psi : \mathcal{D} \to \widetilde{\mathcal{Q}}^{(m)}(\underline{d})$ which is injective. The pullback of the function $\mathfrak{F}_{\underline{\epsilon}}$ by Ψ is given by

$$\Psi^*\mathfrak{F}_{\underline{\epsilon}}(Z) = \exp(-(\frac{|z_1|^2}{\epsilon_1^2} + \dots + \frac{|z_m|^2}{\epsilon_m^2}) - \mathrm{A}(\Psi(Z))).$$

From Theorem 6.5, it suffices to show that

$$\int_{\mathcal{D}} e^{-(|z_1|^2/\epsilon_1^2 + \dots + |z_m|^2/\epsilon_m^2) - \mathrm{A}} d\mu < K\epsilon_1^2 \dots \epsilon_m^2.$$

For this purpose, let us first prove

Lemma 6.6. *The family of coordinates* $(z_1, \dots, z_m)$ *is independent in* $\hat{\mathrm{V}}$.

Proof. Assume that there exists $(\lambda_1, \dots, \lambda_m) \in \mathbb{R}^m$ such that $\lambda_1 Z(e_1) + \dots + \lambda_m Z(e_m) = 0$ for all $Z \in \hat{\mathrm{V}}$. Let $e_{i_k} = \tau(e_k)$, $k = 1, \dots, m$. Since Z satisfies the equations (19), we also have

$$\lambda_1(Z(e_1) + Z(e_{i_1})) + \dots + \lambda_m(Z(e_m) + Z(e_{i_m})) = 0, \ \forall Z \in \hat{\mathrm{V}}. \tag{20}$$

Let $(\hat{M}, \hat{\omega}, \{\hat{\gamma}_1^1, \hat{\gamma}_1^2, \dots, \hat{\gamma}_m^1, \hat{\gamma}_m^2\})$ be the canonical double covering of $\Psi(Z)$, then (e_k, e_{i_k}) corresponds to the pair of saddle connections $(\hat{\gamma}_k^1, \hat{\gamma}_k^2)$ that are exchanged by the involution τ of $\hat{M}$. Note that we have an isomorphism between $\hat{\mathrm{V}}$ and $H^1(M, \hat{S}; \mathbb{C})^- = \{\alpha \in H^1(\hat{M}, \hat{S}; \mathbb{C}) : \ \tau^*\alpha = -\alpha\}$. Condition (20) implies that

$$\alpha(\lambda_1[\hat{\gamma}_1] + \dots + \lambda_m[\hat{\gamma}_m]) = 0, \ \forall \alpha \in H^1(\hat{M}, \hat{S}; \mathbb{C})^-,$$

where $[\hat{\gamma}_k] = [\hat{\gamma}^1_k]+[\hat{\gamma}^2_k]$. Since $H^1(\hat{M},\hat{S};\mathbb{C})^- = \mathrm{Hom}_{\mathbb{R}}(H_1(\hat{M},\hat{S};\mathbb{R})^-,\mathbb{C})$, and $[\hat{\gamma}_k] \in H_1(\hat{M},\hat{S};\mathbb{R})^-,\ k = 1,\dots,m$, it follows that

$$\lambda_1[\hat{\gamma}_1] + \cdots + \lambda_m[\hat{\gamma}_m] = 0 \in H_1(\hat{M},\hat{S};\mathbb{R}).$$

Therefore, we have a contradiction to the hypothesis that the family $\{[\hat{\gamma}_1],\dots,[\hat{\gamma}_m]\}$ is independent in $H_1(\hat{M},\hat{S};\mathbb{R})$. The lemma is then proven. □

The remaining of the proof of the theorem follows the same lines as Theorem 2.3. □

REFERENCES

[EMZ03] A. ESKIN, H. MASUR, A. ZORICH– "Moduli spaces of abelian differentials: The principal boundary, counting problems, and the Siegel-Veech constants", *Publ. Math. Inst. Hautes Études Sci.* **No. 97**, (2003) pp. 61-179.

[EK01] A. ESKIN, and A. OKOUNKOV –"Asymptotics of number of branched coverings of a torus and volume of moduli spaces of holomorphic differentials. *Invent. Math.*, **145:1**, (2001) pp. 59-104.

[EKZ12] A. ESKIN, M. KONTSEVICH, and A. ZORICH – "Sum of Lyapunov exponents of the Hodge bundle with respect to the Teichmüller geodesic flow ", *preprint: arXiv:1112.5872*

[K] M. KONTSEVICH– "Lyapunov exponents and Hodge theory. "The mathematical beauty of physics" " (Saclay, 1996), (in Honor of C. Itzykson) 318-332, *Adv. Ser. Math. Phys., 24. World Sci. Publishing, River Edge*, NJ(1997).

[L04] E. LANNEAU – "Hyperelliptic components of moduli spaces of quadratic differentials with prescribed singularities ", *Comment. Math. Helv.* **79** (2004), pp. 471-501.

[L08] ______– "Connected components of the strata of the moduli spaces of quadratic differentials with prescribed singularities ", *Ann. de l'E.N.S. (4)* **41** (2008), pp. 1-56.

[M82] H. MASUR –"Inteval exchange transformations and measured foliations ", *Annals of Math.* **115** (1982), pp. 169-200.

[MS91] H. MASUR and J. SMILLIE –"Hausdorff dimension of sets of nonergodic foliations ", *Annals of Math. (2)* **134** (1991), no.3, pp. 455-543.

[MT02] H. MASUR and S. TABACHNIKOV – "Rational billards and flat structures ", *Handbook of dynamical systems*, **1A** North-Holland, Amsterdam (2002), pp. 1015-1089.

[MZ08] H. MASUR, and A. ZORICH –"Multiple saddle connections on flat surfaces and the boundary principle of the moduli space of quadratic differentials", *Geom. Funct. Anal.*, **18** (2008), no. 3, pp. 919-987.

[Mc07] C. MCMULLEN – "Dynamics of $\mathrm{SL}_2(\mathbb{R})$ over moduli space in genus two", *Ann. of Math. (2)* **165**, no.2 (2007), pp. 397–456.

[N10] D.-M. NGUYEN – "Triangulation and volume form on moduli spaces of flat surfaces ", *Geom. Funct. Anal.* **20** (2010), no. 1, pp. 192-228.

[N12] D.-M. NGUYEN – "Energy function on moduli spaces of flat surfaces with erasing forest ", *Math. Ann.* **353** (2012) pp. 997-1036.

[Tr91] M. TROYANOV– "Prescribing curvature on compact surfaces with conical singularities", *Trans. Amer. Math. Soc.* **324:2** (1991), pp. 793-821.

[V90] W.A. VEECH –"Moduli spaces of quadratic differentials ", *Journal d'Analyse Math.* **55** (1990), pp. 117-171.

[Z06] A. ZORICH – "Flat surfaces ". In collection *"Frontiers in Number Theory, Physics and Geometry", Vol. 1: On random matrices, zeta functions and dynamical systems*, École de Physique des Houches, France, March 9-21 2003, Springer-Verlag (2006).

IMB BORDEAUX-UNIVERSITÉ BORDEAUX 1
351, COURS DE LA LIBÉRATION
33405 TALENCE CEDEX
FRANCE
E-mail address: `duc-manh.nguyen@math.u-bordeaux1.fr`